arXiv:1304.3022v1 [astro-ph.SR] 10 Apr 2013

Revisiting some physics issues related to the new mass limit for magnetized white dwarfs

Upasana Das and Banibrata Mukhopadhyay
Department of Physics, Indian Institute of Science, Bangalore 560012, India
upasana@physics.iisc.ernet.in , bm@physics.iisc.ernet.in

April 11, 2013

Abstract

We clarify important physics issues related to the recently established new mass limit for magnetized white dwarfs which is significantly super-Chandrasekhar. The issues include, justification of high magnetic field and the corresponding formation of stable white dwarfs, shape of the white dwarfs under consideration, flux freezing, variation of magnetic field and related currents therein. We also attempt to address the observational connection of such highly magnetized white dwarfs.

1 Introduction

Recently, Mukhopadhyay and his collaborators have proposed that highly magnetized white dwarfs could have significantly super-Chandrasekhar masses [1, 2, 3, 4, 5]. Notably, Das & Mukhopadhyay [4] have shown that such white dwarfs have a limiting mass of $2.58M_\odot$. In order to arrive at this result, the authors have made certain assumptions for the convenience of calculation. These are, the choice of constant magnetic field in the central region of white dwarfs, which decreases with density near the surface of the white dwarf, employing the idea proposed by previous authors in the context of strongly magnetized compact objects [6]; the choice of spherical symmetry for strongly magnetized white dwarfs; separate central and surface flux conservation in magnetized white dwarfs as they evolve (e.g., due to accretion), etcetera.

Now, in the present report, we plan to justify the above assumptions in a greater detail, in the light of various related physics issues. This is particularly important, as over the last few months several friends, colleagues and critics have raised certain questions about our proposal of significantly super-Chandrasekhar white dwarfs and a new mass limit of white dwarfs. This report is by and large a suitable compilation of our responses to about a dozen referee reports, which we have successfully tackled during this series of works starting with the one that appeared in 2012 [1].

2 Equipartition magnetic field for super-Chandrasekhar white dwarfs

In this section, we would like to clarify that for the super-Chandrasekhar white dwarfs considered in our works, the equipartition magnetic field (B) is no longer same as that for weakly magnetized white dwarfs, but much higher.

One can estimate the equipartition field of a star by recalling the scalar virial theorem [7]. By equating the magnetic and the gravitational potential energies of a spherical star of mass M and radius R, i.e., $(4\pi R^3/3)(B_{\rm equi}^2/8\pi) \sim GM^2/R$, one arrives at the maximum possible equipartition magnetic field $B_{\rm equi} \sim 2\times 10^8 (M/M_\odot)(R/R_\odot)^{-2}$ G, where $M_\odot$ and $R_\odot$ are the mass and radius of Sun respectively and G is Newton's gravitation constant. For a typical white dwarf having radius $\sim 10^4$ km and mass $1.4M_\odot$, this turns out to be $\sim 10^{12}$ G. However, for a super-Chandrasekhar white dwarf having, for example, mass $2.58M_\odot$ and radius ~ 70 km (see, e.g., Figure 1 of [5]), $B_{\rm equi}$ is $\sim 5\times 10^{16}$ G $\gg 10^{12}$ G. It is to be noted, however, that this is but an average value and hence is less than the value of the field considered to model the central region of the above white dwarf, and larger than the value at the surface. In fact, if the central region is defined to be the region of radius $R_{int} = R/10$ which demarcates a zone up to which the central magnetic field of the white dwarf remains (almost) constant [5], then the above argument yields $B_{\rm equi} \sim 5\times 10^{18}$ G in this zone. This is very well consistent with the central magnetic field B_{int} ($= 8.8\times 10^{17}$ G) used to model the white dwarf considered above. Hence, although we are considering super-critical (central) magnetic fields ($B > B_c = 4.414\times 10^{13}$ G), which cause Landau quantization to significantly alter the equation of state, we are *not* actually considering super-equipartition fields. Note that the value of $B_{\rm equi}$ itself has increased for a super-Chandrasekhar white dwarf, since with the increase in the magnetic field strength, the white dwarfs start becoming more compact in size (see, e.g., Figures 4(b), (c) and §IV.B of [2]). We would like to point out here that, indeed, Ostriker and Hartwick [8] constructed models of white dwarfs with $B \gtrsim 10^{12}$ G at the center but with a much smaller field at the surface. Thus, the concept of high interior magnetic fields in white dwarfs, although difficult to verify observationally, are not implausible.

3 Justification of constant central magnetic field

In our works, we have modeled the central region of the white dwarf to have a strong constant magnetic field (several orders of magnitude larger than the surface field). This can be justified as follows.

From Maxwell's equations in a steady state, we have

$$\nabla \times \mathbf{B} = \frac{4\pi}{c}\mathbf{j}, \tag{1}$$

where $\mathbf{j}$ is the current density. Now, in the presence of a very strong central magnetic field, $B \gg B_c$ (as is the case for a significantly super-Chandrasekhar white dwarf), the electron degenerate matter will not only be Landau quantized but also the electrons will occupy mostly the ground Landau level. Thus the electrons are not expected to move and hence practically there will be no current, i.e., $\mathbf{j} = 0$. Therefore, from the above equation, we obtain that there is no spatial variation of B, i.e., the magnetic field is constant. Since it is this central magnetic field which is primarily responsible for the mass to exceed the Chandrasekhar limit (see, e.g., §IV.C of [2]), one can consider the hydrostatic equilibrium condition in order to solve for the structure of the magnetized white dwarfs, at least in the central region, for obtaining super-Chandrasekhar white dwarfs.

4 Stable equilibrium in presence of strong magnetic field

One might wonder whether magnetic energy could really dominate over internal energy without magnetic forces significantly altering the equilibrium. Since the central region of the white dwarf in our works is modeled to have a constant magnetic field, the magnetic forces vanish therein. Nevertheless, the field must drop off by the time the surface is reached and hence the field gradients (and therefore magnetic forces) could exist primarily in the lower density outer regions, which are, however, less important for the super-Chandrasekhar mass calculation (see, e.g., Figure 5 and §IV.C of [2]). We demonstrate in this section that gravitational forces in the low density regions would still dominate over the magnetic forces making it possible for the white dwarf to attain a stable equilibrium.

Let us consider a super-Chandrasekhar white dwarf having a mass $2.39M_\odot$ (which is less than the mass limit of $2.58M_\odot$), radius 9×10^7 cm and a central magnetic field $B_{int} = 4.414\times10^{15}$ G (this is the white dwarf corresponding to the maximum mass point on the dotted line in Figure 2 of [4]). Figure 1(a) shows the variation of density as a function of radius inside the above white dwarf having a central density $\rho_c = 4.14\times10^9$ gm/cc. Figures 1(b) and 1(c) show the variation of matter pressure and mass as functions of radius. If we consider a radius in the low density region of this white dwarf $R' = 7 \times 10^7$ cm, then from the figures we see that the density at this radius is $\rho(R') = 1.46 \times 10^9$ gm/cc, the mass within this radius is $M(R') = 1.73M_\odot$ and the matter pressure is $P(R') = 8.83 \times 10^{26}$ ergs/cc. Adopting the description of the variation of magnetic field with density in a compact star given by Equation 5 of Bandyopadhyay et al. [6], we show a plausible profile of the field in Figure 1(d) below, using suitable values for the parameters $\beta = 0.001$ and $\gamma = 6$ (note, it is not the same as the polytropic index) and the values of central and surface magnetic fields. We note that the profile assures almost constant B in the central region of the white dwarf and then B drops off near the surface. Since it is the central field which is responsible for contributing significant mass to the white dwarf, such a profile would produce very similar results to those we have already obtained. From Figure 1(d) we obtain that the magnetic field at $\rho(R')$ is $B(R') = 1.6 \times 10^{14}$ G.

Now, in the presence of a varying magnetic field the white dwarf must obey the condition of magnetostatic equilibrium:

$$\frac{1}{\rho}\frac{d}{dr}\left(P + \frac{B^2}{8\pi}\right) = -\frac{GM}{r^2} + \left[\frac{\vec{B}\cdot\nabla\vec{B}}{4\pi\rho}\right]_r, \tag{2}$$

where, r is the radial distance from the center of the white dwarf, P and ρ are the pressure and density of the electron degenerate matter respectively. We now present order of magnitude estimates of the various terms of Equation (2) in the low density region of the white dwarf, where the gradient of magnetic field is non-zero. The orders of gravitational, pressure and magnetic forces per unit mass, at R', are $GM(R')/R'^2$, $P(R')/(\rho(R')R')$ and



Figure 1: (a) Density, (b) pressure and (c) mass, as functions of radius, within a white dwarf of mass $2.39M_\odot$ and central magnetic field 4.414×10^{15} G. (d) Variation of magnetic field with density inside the same white dwarf. ρ_0 is a normalization density with a value of 8×10^8 gm/cc. R is in units of cm, ρ in gm/cc, P in ergs/cc, M in $M_\odot$ and B in G.

$B(R')^2/(8\pi R'\rho(R'))$ respectively. Hence at R' the gravitational, pressure and magnetic forces per unit mass are of the order 4.71×10^{10} dyne/gm, 8.64×10^9 dyne/gm and 9.96×10^9 dyne/gm respectively. Thus we see that the gravitational force still dominates over the magnetic force (and combined rest of the forces) in the low density region and hence it is possible to construct a stable equilibrium solution. Note that, when exact gradients are calculated in reality, gravitational force will exactly balance the combined forces due to matter pressure and magnetic field.

5 Degenerate matter density stronger than magnetic density

One can also make stability arguments by computing the magnetic density ($\rho_B = B^2/8\pi c^2$) and the matter density (ρ) inside the white dwarf shown in Figure 1 and then their effects on the gravitational energy density. Now, the mass of the white dwarf depends directly on the matter density and the gravitational energy density is in turn proportional to the mass. Hence, higher the matter density, higher is the gravitational energy density. The magnetic density and the matter density at the center of the white dwarf of mass $2.39 M_\odot$ are 8.6×10^8 gm/cc and 4.14×10^9 gm/cc respectively. Thus the matter density is about an order of magnitude larger than the magnetic density. The magnetic field remains constant in the central region (and hence so does ρ_B) up to a matter density $\sim 3 \times 10^9$ gm/cc (see Figure 1(d)), but still we note that matter density is greater than $\rho_B = 8.6 \times 10^8$ gm/cc. As the density decreases further, the magnetic field starts falling off near the surface of the white dwarf. For example, the magnetic densities $\rho_B = 2.1 \times 10^8$ gm/cc, 3.5×10^6 gm/cc and 7.8×10^3 gm/cc correspond to the matter densities $\rho = 2.4 \times 10^9$ gm/cc, 1.6×10^9 gm/cc and 9.6×10^8 gm/cc, respectively. Thus we see that, ρ_B decreases much faster than ρ as the surface of the white dwarf is approached and hence a stable equilibrium can be maintained throughout the white dwarf and ρ_B can be ignored to compute the mass of white dwarfs. Note that, if in the central region the contribution from ρ_B is included in order to compute the mass of the white dwarfs, then this may slightly change the mass estimate and probably even increase it.

6 Stability against magnetic buoyancy

A question that may arise is the issue of buoyancy related instabilities within the strongly magnetized super-Chandrasekhar white dwarfs. If a blob of gas is displaced adiabatically from an inner radius to an outer radius in the central region of the white dwarf, then the condition for stability is given by:

$$\rho' = \left(\frac{K_1}{K_2} \rho_1^{\Gamma_1} \right)^{1/\Gamma_2} > \rho_2, \tag{3}$$

where ρ' is the matter density inside the blob when it is at the outer radius, ρ_1, Γ_1 and K_1 are the matter density, adiabatic index and a dimensional constant related to the adiabatic equation of state respectively of the surrounding medium at the inner radius and ρ_2, Γ_2 and K_2 are the same quantities at the outer radius. Note that, since the magnetic field is constant in the central region of the white dwarf, the magnetic pressure



Figure 2: Variation of the difference in densities inside and outside a displaced blob of gas as a function of its surrounding density. Density is in units of gm/cc.

terms do not appear in the above relation. The above criterion means that the blob at the outer radius must be heavier than the surrounding (i.e., $\rho' - \rho_2 > 0$) so that it is pulled inwards by gravity to its original position, making the system stable. If the blob is lighter than its surrounding (i.e., $\rho' - \rho_2 < 0$) then the buoyant force displaces the blob further outwards, making the system unstable.

We apply this argument to a white dwarf having central density $\sim 1.4 \times 10^9$ gm/cc; the equation of state of the underlying degenerate gas being described by the parameters listed in Table II of [2]. In Figure 2 we show the variation of $\rho' - \rho_2$ with ρ_2 inside this white dwarf in a few discrete radial zones. We clearly see that $\rho' - \rho_2$ is everywhere positive, establishing that the white dwarf is stable against buoyancy related instabilities. Away from the central region, the magnetic field decreases towards the surface (as discussed previously), so the magnetic pressure at the outer radius will be less than that at the inner radius. Taking this difference into account will only increase the density of the blob, making it even more heavier than the surrounding, further strengthening the argument for a stable system.

7 Choice of spherical symmetry

Order of magnitude estimates obtained in §4 argue that the overall gravitational energy in the white dwarfs under consideration dominates over the combined total energy that arises from matter and magnetic field. This justifies our choice of spherical symmetry.

However, if the magnetic field is very strong, then the pressure may become anisotropic [9]. Following Bocquet et al. [10], the combined fluid-magnetic medium then may develop a tension at such a high magnetic field. Bocquet et al. [10] observed that as a result of this magnetic tension, the star displays a pinch across the symmetry axis and assumes a flattened shape. It is possible that a similar effect could occur in highly magnetized white dwarfs. In that case, the magnetic tension would deform the magnetized white dwarf along the direction of the magnetic field and possibly turn it into a kind of oblate spheroid. Consequently, the spherically symmetric hydrostatic equilibrium equation would have to be modified due to the presence of the strong magnetic field. As stated by Paulucci et al. [11], one has to generalize the Tolman-Oppenheimer-Volkoff equation to take into account the anisotropy in the pressure, but to the best of our knowledge this has not been achieved so far, as is also mentioned in the same paper itself. We propose to carry out such calculations for highly magnetized white dwarfs in a future work. However, in the previous work [2], we have initiated the understanding of the effect of deviation from spherical symmetry due to the magnetic field, if any, in the Appendix. Interestingly we observe that, flattening effects due to magnetic field renders super-Chandrasekhar white dwarfs even at relatively weaker magnetic fields which are more probable in nature.

Important point to note, however, is that, for the strongly magnetized white dwarf having the new limiting mass [4], the above geometrical factors do not matter, since the corresponding radius is negligible (as is also the case for the Chandrasekhar limit).

8 Difficulty behind direct observation of strongly magnetized super-Chandrasekhar white dwarfs

Recent observations of several peculiar over-luminous type Ia supernovae - SN 2006gz, SN 2007if, SN 2009dc, SN 2003fg - seem to suggest super-Chandrasekhar-mass white dwarfs with masses up to $2.1 - 2.8 M_\odot$ as their most likely progenitors [12, 13]. The strongly magnetized super-Chandrasekhar white dwarfs proposed by Das & Mukhopadhyay [2, 4] and Das, Mukhopadhyay & Rao [5] can explain these supernova explosions, which is their indirect evidence. However, there has been no direct detection of such white dwarfs so far. We try to explore the reason behind this in this section.

Ordinary sub-Chandrasekhar white dwarfs are located in the lower-left portion of the Hertzsprung-Russell (H-R) diagram (plot showing the relationship between the luminosities of stars and their effective temperatures), indicating that they are low luminosity hot objects. If the surface of the white dwarf is considered to radiate like a black body, then its luminosity is given by $L = 4\pi R^2 \sigma_B T_{eff}^4$, where σ_B is the Stefan-Boltzmann constant and T_{eff} the effective surface temperature of the white dwarf. Since $L \propto R^2$, ordinary white dwarfs, having radius $\sim 10^4$ km, are bright enough to feature in the H-R diagram. Neutron stars on the other hand, having radius 10 km, are too small (and hence too dim) and would lie far below the white dwarfs in the H-R diagram, and hence do not not appear in it altogether. Note that, neutron stars are mostly detected as pulsars, due to the emissions of regular pulses. The strongly magnetized super-Chandrasekhar white dwarfs in our works, having radius $\lesssim 100$ km, are more closer to neutron stars than conventional white dwarfs. Thus, such (isolated) white dwarfs would also *not* appear in the H-R diagram, making them virtually impossible to be detected directly based on current observational techniques.

9 Separate flux conservation in the central and surface regions

Das, Mukhopadhyay & Rao [5] have discussed a possible evolutionary scenario from weakly magnetized, sub-Chandrasekhar, accreting white dwarfs to strongly magnetized, super-Chandrasekhar white dwarfs, which eventually explode to give rise to the peculiar type Ia supernovae. Irrespective of how the interior magnetic field of the white dwarf is modeled, the only constraint on the above evolution is the magnetic flux conservation between the initial and final white dwarfs, which controls/relates the magnetic field profiles of the white dwarfs, if one of them (say the initial one) is known. Additionally, the authors have imposed that the magnetic flux is conserved separately in the central and surface regions of the white dwarfs. This is justified as follows.

In §2 of [5], the authors have defined a radius R_{int} which demarcates a zone up to which the central magnetic field of the white dwarf remains (almost) constant which is larger than that at the surface. This is quite a plausible assumption, since the original star which collapses to form a white dwarf might have a very large interior magnetic field compared to that observed on its surface. For example, Parker [14] estimated that a magnetic field strength as high as $\sim 10^7$ G can not be ruled out for the Sun's interior. Gough & McIntyre [15] also discussed the inevitability of a strong magnetic field in the Sun's interior and its influence to the extent of a transitional layer, called the tachocline. Thus, it is quite likely that the values of the central and surface magnetic fluxes (in the initial star and hence in the white dwarf) and the underlying electric currents might be significantly different. Therefore, the size of the central (and then the surface) region of a white dwarf is controlled by the size of the central region of the original star. Since it is likely that there is a gradient of magnetic field from the center to the surface of the star, the radial dependence of B could be of the form $B \propto 1/r^{\zeta}$, where ζ is positive. However, the above field profile will blow up at $r = 0$ which is unfeasible and hence B is expected to be very large and to remain constant in the central region. This restricts the electrons to move around in the central region of the final highly magnetized white dwarfs, as only the lower Landau level(s) is/are available for them, justifying practically no current therein. Nevertheless, as B decreases towards the surface, higher Landau levels become available and hence the electrons can produce current, maintaining a gradient of the magnetic field up to the surface.

All the above arguments justify the assumption of the separate conservation of the surface and central magnetic fluxes, which also suggest the presence of a core-envelope boundary.

10 Correlation between the observed peculiar type Ia supernovae rate and the occurrence of highly magnetized super-Chandrasekhar white dwarfs

Identification of the progenitors of type Ia supernovae is still an ongoing research area. In [5] we have proposed that if the progenitors are single degenerate white dwarfs accreting matter from a normal companion, then the peculiar supernovae originate from super-Chandrasekhar white dwarfs with extremely high magnetic fields.

The observed rate of over-luminous, peculiar type Ia supernovae is about 1-2% of the normal type Ia [16]. This is consistent with the fact that about 25% of all cataclysmic variables (CVs) are magnetic CVs and we expect a small fraction (say 10%) of them to be super-magnetic. These high magnetic CVs (2.5% of all CVs based on the above assumption) would behave like normal magnetic CVs during most of their life time, till their mass remains sub-Chandrasekhar. However, they would be seen as peculiar objects only during a small fraction of their life time, when the surface magnetic field becomes very high ($> 10^{11}$ G) and the mass is super-Chandrasekhar ($> 1.44M_\odot$), which should happen because of accretion. We have discussed this issue with constant and varying accretion rates in §3 of [5].

For a constant accretion rate of $2.5\times10^{-9}M_\odot\mathrm{yr}^{-1}$, which leads to a white dwarf having final mass $2.33M_\odot$ (shown by the dashed line in Figure 3(a) of [5]), the white dwarf spends about 0.4 fraction of its lifetime with super-Chandrasekhar mass. For varying accretion rate, modeled by Equation 4 of [5] with $\alpha = 2$ (solid line in Figure 3(b) of [5]), the white dwarf spends about 0.05 fraction of its lifetime with super-Chandrasekhar mass. Now, typically observed stably accreting intermediate polars have accretion rates $\sim 10^{-9}M_\odot\mathrm{yr}^{-1}$ [17]. However, an accretion rate $\gtrsim 10^{-7}M_\odot\mathrm{yr}^{-1}$ is required for the stable burning of the accreted hydrogen and helium in (single degenerate) type Ia supernovae progenitors [18], which is indeed attained by the above model with varying accretion rate, starting with an accretion rate similar to that of intermediate polars. Thus, the varying accretion rate is a more likely scenario, which gives us the estimate that about 0.12 % of CVs would be peculiar high magnetic CVs. Since they are accreting at a high rate, they will be visible as bright X-ray objects, rather than white dwarfs in optical data bases like SDSS.

11 Summary

We have reestablished the subtle physics issues related to highly magnetized super-Chandrasekhar white dwarfs. We have shown that such white dwarfs are indeed possible, provided certain obvious physics constraints are satisfied, explained in this report. This further strengthens the argument for the existence of such super-Chandrasekhar white dwarfs.

Acknowledgments

The authors would like to thank A. R. Rao of TIFR, Diptiman Sen, Tarun Deep Saini and Prateek Sharma of IISc for discussion and encouragement. This work is partly supported by ISRO project Grant No. ISRO/RES/2/367/10-11. U.D. thanks CSIR, India for financial support.

References

[1] A. Kundu and B. Mukhopadhyay, Mod. Phys. Lett. A **27**, 1250084 (2012)

[2] U. Das and B. Mukhopadhyay, Phys. Rev. D **86**, 042001 (2012)

[3] U. Das and B. Mukhopadhyay, Int. J. Mod. Phys. D **21**, 1242001 (2012)

[4] U. Das and B. Mukhopadhyay, Phys. Rev. Lett. **110**, 071102 (2013)

[5] U. Das, B. Mukhopadhyay and A.R. Rao, ApJ **767**, L14 (2013)

[6] D. Bandyopadhyay, S. Chakrabarty and S. Pal, Phys. Rev. Lett. **79**, 2176 (1997)

[7] S.L. Shapiro and S.A. Teukolsky, *Black Holes, White Dwarfs and Neutron Stars: The Physics of Compact Objects* (John Wiley and Sons, New York, 1983).

[8] J.P. Ostriker and F.D.A. Hartwick, ApJ **153**, 797 (1968)

[9] M. Sinha, B. Mukhopadhyay and A. Sedrakian, Nucl. Phys. A **898**, 43 (2013).

[10] M. Bocquet, S. Bonazzola, E. Gourgoulhon and J. Novak, A&A **301** 757 (1995)

[11] L. Paulucci, E.J. Ferrer, V. de la Incera and J.E. Horvath, Phys. Rev. D **83**, 043009 (2011)

[12] D.A. Howell, et al., Nature **443**, 308 (2006)

[13] R.A. Scalzo, et al., ApJ **713**, 1073 (2010)

[14] E.N. Parker, *Cosmical magnetic fields: Their origin and their activity* (Oxford University Press, 1979)

[15] D.O. Gough and M.E. McIntyre, Nature **394**, 755 (1998)

[16] R.A. Scalzo, et al., ApJ **757**, 12 (2012)

[17] B. Warner, *Cataclysmic Variable Stars* (Cambridge University Press, 1995)

[18] A. Cumming, MNRAS, **333**, 589 (2002)